

---

# *Overview of **BTeV** Physics, the Components and the Requirements*

# The Physics: General


## Mysteries

## Solutions: New Physics

Dark  
Matter


Dominance  
of Matter over  
Antimatter

Dark  
Energy



# The Physics: More Specific

- CP Violation: Particles behave differently than antiparticles
  - Demonstrated in B decays by BaBar & Belle (one  $\angle$  measured,  $\beta$ )
  - But there are 4 different angles to determine
  - New Physics can show up as inconsistencies between/among CP measurements and other quantities
- Rare Decays




- New Particles can appear in the loop & interfere

# Project Scope

WBS 1.0

WBS 3.0



WBS 4.0

**BTeV Project**

WBS 2.0



## Requirements on C0 IR

---

- Peak Luminosity  $\sim 2 \times 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$  ( $\beta^* < 50 \text{ cm}$ )
- Interoperability: Must allow for operation at C0 or at B0 & D0 simultaneously
- Non-interference with BTeV detector – last quadrupole closest to collision point is 5 m further away than in CDF or D0
- Schedule: Must be ready by shutdown in middle of 2009

# ~~BTeV~~ C0 Requirements on C0 Outfitting (*WBS 3.0*)

---

- Building already exists
- We need to
  - Provide the architectural, structural, mechanical and electrical work for the BTeV detector (WBS 1.0).
  - Provide the modifications to the C-0 Service Building and primary power for the Interaction Region (WBS 2.0).

**Belarussian State-** D .Drobychev,  
A. Lobko, A. Lopatrik, R. Zouversky

**UC Davis** - P. Yager

**Univ. of Colorado at Boulder**

J. Cumalat, P. Rankin, K. Stenson

**Fermi National Lab**

J. Appel, E. Barsotti, C. Brown,  
J. Butler, H. Cheung, D. Christian,  
S. Cihangir, M. Fischler,  
I. Gaines, P. Garbincius, L. Garren,  
E. Gottschalk, A. Hahn, G. Jackson,  
P. Kasper, P. Kasper, R. Kutschke,  
S. W. Kwan, P. Lebrun, P. McBride,  
J. Slaughter, M. Votava, M. Wang,  
J. Yarba

**Univ. of Florida at Gainesville**  
P. Avery

**University of Houston –**

A. Daniel, K. Lau, M. Ispiryan,  
B. W. Mayes, V. Rodriguez,  
S. Subramania, G. Xu

**Illinois Institute of Technology**

R. Burnstein, D. Kaplan,  
L. Lederman, H. Rubin, C. White

**Univ. of Illinois-** M. Haney, D.  
Kim, M. Selen, V. Simatis, J. Wiss

**Univ. of Insubria in Como-**  
P. Ratcliffe, M. Rovere

**INFN - Frascati-** M. Bertani, L.  
Benussi, S. Bianco, M. Caponero, D.  
Collona, F. Fabri, F. Di Falco, F.  
Felli, M. Giardoni, A. La Monaca,  
E. Pace, M. Pallota, A. Paolozzi, S.  
Tomassini

**INFN - Milano** – G. Alimonti,  
P’Dangelo, M. Dinardo, L. Edera, S.  
Erba, D. Lunesu, S. Magni, D.  
Menasce, L. Moroni, D. Pedrini, S.  
Sala, L. Uplegger

**INFN - Pavia** - G. Boca, G.  
Cossali, G. Liguori, F. Manfredi, M.  
Maghisoni, L. Ratti, V. Re, M.  
Santini, V. Speviali, P. Torre, G.  
Traversi

**IHEP Protvino, Russia** - A.  
Derevschikov, Y. Goncharenko, V.  
Khodyrev, V. Kravtsov, A.  
Meschanin, V. Mochalov,  
D. Morozov, L. Nogach, P.  
Semenov K. Shestermanov,  
L. Soloviev, A. Uzunian, A. Vasiliev

**University of Iowa**

C. Newsom, & R. Braunger

**University of Minnesota**

J. Hietala, Y. Kubota, B. Lang,  
R. Poling, A. Smith

**Nanjing Univ. (China)-**

T. Y. Chen, D. Gao, S. Du,  
M. Qi, B. P. Zhang, Z. Xi  
Xang, J. W. Zhao

**New Mexico State -**

V. Papavassiliou

**Northwestern Univ. -**

J. Rosen

**Ohio State University-**

K. Honscheid, & H. Kagan

**Univ. of Pennsylvania**

W. Selove

**Univ. of Puerto Rico**

A. Lopez, H. Mendez, J.  
Ramirez, W. Xiong

**Univ. of Science & Tech.**

**of China** - G. Datao, L. Hao,  
Ge Jin, L. Tiankuan, T. Yang,  
& X. Q. Yu

**Shandong Univ. (China)-**

C. F. Feng, Yu Fu, Mao He,  
J. Y. Li, L. Xue, N. Zhang, &  
X. Y. Zhang

**Southern Methodist –**

T. Coan, M. Hosack

**Syracuse University-**

M. Artuso, C. Boulahouache,  
S. Blusk, J. Butt, O.

Dorjkhaidav, J. Haynes, N.

Menaa, R. Mountain,

H. Muramatsu, R. Nandakumar,

L. Redjimi, R. Sia,

T. Skwarnicki, S. Stone, J. C.

Wang, K. Zhang

**Univ. of Tennessee**

T. Handler, R. Mitchell

**Vanderbilt University**

W. Johns, P. Sheldon,

E. Vaandering, & M. Webster

**University of Virginia** M.

Arenton, S. Conetti, B. Cox, A.

Ledovskoy, H. Powell, M.

Ronquest, D. Smith, B.

Stephens, Z. Zhe

**Wayne State University**

G. Bonvicini, D. Cinabro,

A. Schreiner

**University of Wisconsin**

M. Sheaff



**York University** - S. Menary

# $\frac{BTeV}{C0}$ Characteristics of hadronic $b$ production

$$p\bar{p} \rightarrow b\bar{b} + X$$

The higher momentum  $b$ 's are at larger  $\eta$ 's

$b$  production peaks at large angles with large  $b\bar{b}$  correlation





# Requirements: General

---

- Intimately tied to Physics Goals
- In general, within the acceptance of the spectrometer (10 – 300 mr with respect to beam) we need to:
  - Detect charged tracks & measure their 3-momenta
  - Measure the point of origin of the charged tracks (vertices)
  - Detect neutrals & measure their 3-momenta
  - Reveal the identity of charged tracks (e,  $\mu$ ,  $\pi$ , K, p)
  - Trigger & acquire the data (DAQ)
- Detector we designed meets the requirements


## Basics Reasons for the Requirements

- B's (& D's) are long lived,  $\sim 1.5$  ps, so if they are moving with reasonable velocity they go  $\sim 3$  mm before they decay. This allows us to Trigger on the the presence of a B decay (*detached vertex*).
- B's are produced in pairs  $p\bar{p} \rightarrow b\bar{b} + X$ , and for many crucial measurements we must detect one b fully and some parts of the other: “flavor tagging”
- Physics states of great interest now are varied and contain both charged modes and neutrals,  $B_d$  &  $B_s$

## More Basic Reasons


- Many modes contain  $\gamma$ ,  $\pi^0$  &  $\eta$ , so need excellent electromagnetic calorimetry
- $B_s$  oscillations are fast, so need excellent time resolution  $\sim < 50$  fs, compared to  $\sim 1500$  fs lifetime. Also very useful to reduce backgrounds in reconstructed states
- Physics Backgrounds from  $\pi \Leftrightarrow K$  can be lethal
  - $B_s \rightarrow D_s \pi^-$  is 15X  $B_s \rightarrow D_s K^-$
  - $B^0 \rightarrow K^* \pi \rightarrow K^\mp \pi^\pm \pi^0$  is 2X  $B^0 \rightarrow \rho \pi \rightarrow \pi^+ \pi^- \pi^0$
  - So excellent charged hadron identification is a must

# $\frac{BTeV}{C0}$ The BTeV detector in the C0 collision hall



# The BTeV Detector



## BTeV Detector Layout



# BTeV Co Fundamentals: Decay Time Resolution

- Excellent decay time resolution
  - Reduces background
  - Allows detached vertex trigger
- The average decay distance and the uncertainty in the average decay distance are functions of B momentum:

$$\begin{aligned}\langle L \rangle &= \gamma \beta c \tau_B \\ &= 480 \mu\text{m} \times p_B / m_B\end{aligned}$$




CDF/D0  
region →

↖ LHC-b  
region

# Pixels (WBS 1.2)


- Pixel – working systems studied in beams, including “almost” final electronics
- Full mechanical design done and being tested
- Pixels are inside of beam pipe in machine vacuum – OK with accelerator provided the outgassing rate is below specified limits (review document linked to Review web page)



- Full GEANT has multiple scattering, bremsstrahlung, pair conversions, hadronic interactions and decays in flight; smears hits and refits the tracks using “Kalman Filter.” No pattern recognition (except for trigger). However, we do not expect large pattern recognition problems

- This track density is 3x higher than what is expected in BTeV!


- ◆ Detailed studies of efficiency and rejection for up to an average of six interactions/crossing





# <sup>BTeV</sup><sub>Co</sub> Pixel Trigger Overview (WBS 1.8)

◆ Idea: find primary vertices & detached tracks from b or c decays



- Pixel hits from 3 stations are sent to an FPGA tracker that matches “interior” and “exterior track hits
- Interior and exterior triplets are sent to a farm of DSPs to complete the pattern recognition:
  - interior/exterior triplet matcher
  - fake-track removal


# Trigger Performance

- For a requirement of at least 2 tracks detached by more than  $4\sigma$ , we trigger on only 1% of the beam crossings and achieve the following efficiencies for these states at Level I:


State	efficiency(%)	state	efficiency(%)
$B \rightarrow \pi^+ \pi^-$	55	$B^0 \rightarrow K^+ \pi^-$	54
$B_s \rightarrow D_s K$	70	$B^0 \rightarrow J/\psi K_s$	50
$B^- \rightarrow D^0 K^-$	60	$B_s \rightarrow J/\psi K^*$	69
$B^- \rightarrow K_s \pi^-$	40	$B^0 \rightarrow K^* \gamma$	40

@ 2 int/crossing

- Straws (*WBS 1.6*):  
prototype undergoing  
tests, uses Atlas design  
as basis




- Silicon Strips  
(*WBS 1.7*): simple  
single sided design,  
mechanics done.






# $BTeV$ Co RICH (WBS 1.3) : Two Systems

- Gas + Mirror + MAPMT to identify b decay products
- Liquid + PMT's to help with flavor tagging of b's (p/K separation for  $p < 9 \text{ GeV}/c$ )
- Excellent particle id. distinguishes BTeV from “Central pp Detectors”




# $BTeV$ Co EM cal (WBS 1.4) using $PbWO_4$ Crystals

- GEANT simulation of  $B^0 \rightarrow K^* \gamma$ , for BTeV & CLEO
- Isolation & shower shape cuts on both



Radius (cm)






CLEO  
barrel  
 $\epsilon = 89\%$

# $B^0 \rightarrow \rho \pi$

Based  $9.9 \times 10^6$  bkgrnd events

$B^0 \rightarrow \rho^+ \pi^-$  S/B = 4.1

$B^0 \rightarrow \rho^0 \pi^0$  S/B = 0.3



# Muon System (*WBS 1.5*)

- Used to check detached vertex trigger by having an independent di-muon trigger
- Also used for  $\mu$  id
- Tested in beams
- Robust design: stainless steel tubes





# Kinds of Requirements

---

- One set of requirements is based on the physics performance we want the detector to provide
- A second set is internal to the detector subsystem of interest and tells how each individual piece needs to perform (i. e. the efficiencies of PM tubes, or noise on electronics)
- I will concentrate on the first set here



- Luminosity up to  $2 \times 10^{32} \text{ cm}^{-2}\text{s}^{-1}$
- Mean number of interactions per crossing of 6 (thus allowing for 396 ns bunch spacing)
- Time between bunches  $< 100 \text{ ns}$  (thus allowing for 132 ns bunch spacing)
- Radiation Resistance for at least 10 years on all detector components

# High Level Requirements

---

## ■ Charged Tracks

- Angular acceptance: 10 - 300 mr
- $p > 3 \text{ GeV}/c$
- Tracking efficiency  $> 98\%$
- Mass resolution  $< 50 \text{ MeV}/c$
- Primary vertex resolution (along beam)  $< 100 \mu\text{m}$

## ■ Trigger efficiency & rejection

- $\epsilon > 50 \%$  for all B decays with  $\geq 2$  charged tracks
- $\epsilon > 20 \%$  for all B decays with 1 charged track
- Trigger rejection  $> 98\%$  on light quark events (Level I), and  $99.9\%$  at Level III with only a 10% further loss in b efficiency
- Maximum data rate to archival storage  $< 200 \text{ Mbyte/sec}$

# <sup>BTeV</sup> Co Hadron & Lepton Identification

---

- $\pi/K$  separation  $\geq 4\sigma$  for momenta 3 - 70 GeV/c
- $p/K$  separation  $\geq 3\sigma$  for momenta 3 - 70 GeV/c
  - These allow for  $\pi/e$  &  $\pi/\mu$  separation at  $4\sigma$  level up to  $\sim 23$  and  $\sim 17$  GeV/c, respectively
- positive  $\mu$  identification from 5 - 100 GeV/c with a fake rate  $< 10^{-3}$  and an independent momentum determination with resolution

$$\frac{\sigma_p}{p} = 19\% \oplus 0.6\% \times p$$

# ~~BTeV~~ Co Electromagnetic Calorimeter (WBS 1.4)

- Radius up to 160 cm ~220 mr, with hole for beam ~10 mr
- Range  $E > 1$  GeV
- Energy resolution

$$\frac{\sigma_E}{E} < \frac{2\%}{\sqrt{E}} \oplus 1\%$$

- Position resolution

$$\sigma_x < \frac{4 \text{ mm}}{\sqrt{E}} \oplus 1 \text{ mm}$$

# BTeV co Physics Reach (CKM) in $10^7$ s

Reaction	$\mathcal{B}(B)(\times 10^{-6})$	# of Events	S/B	Parameter	Error or (Value)
$B^0 \rightarrow \pi^+ \pi^-$	4.5	14,600	3	Asymmetry	0.030
$B_s \rightarrow D_s K^-$	300	7500	7	$\gamma$	$8^\circ$
$B^0 \rightarrow J/\psi K_S \quad J/\psi \rightarrow l^+ l^-$	445	168,000	10	$\sin(2\beta)$	0.017
$B_s \rightarrow D_s \pi^-$	3000	59,000	3	$x_s$	(75)
$B^- \rightarrow D^0 (K^+ \pi^-) K^-$	0.17	170	1		
$B^- \rightarrow D^0 (K^+ K^-) K^-$	1.1	1,000	>10	$\gamma$	$13^\circ$
$B^- \rightarrow K_S \pi^-$	12.1	4,600	1		$< 4^\circ +$
$B^0 \rightarrow K^+ \pi^-$	18.8	62,100	20	$\gamma$	theory errors
$B^0 \rightarrow \rho^+ \pi^-$	28	5,400	4.1		
$B^0 \rightarrow \rho^0 \pi^0$	5	780	0.3	$\alpha$	$\sim 4^\circ$
$B_s \rightarrow J/\psi \eta, \quad J/\psi \rightarrow l^+ l^-$	330	2,800	15		
$B_s \rightarrow J/\psi \eta'$	670	9,800	30	$\sin(2\chi)$	0.024

# Endorsements & Schedule

- BTeV was included as a near term priority in the category of “Highest Scientific Importance and Near-term Readiness for Construction,” in the “Facilities for the Future of Science: A Twenty-year Outlook” report of the Office of Science.
- Based on our physics sensitivities, and **implementation in 2009** a HEPAP subpanel wrote: “P5 supports the construction of BTeV as an important project in the world-wide quark flavor physics area. Subject to constraints within the HEP budget, **we strongly recommend an earlier BTeV construction profile** and enhanced C0 optics”
- *Presidents FY2005 Budget Request: “The BTeV experiment will have scientific competition from a dedicated B physics experiment at the CERN LHC, so timely completion of BTeV is important.”*